? Eyster

wayy

9-75 u

activated

"Absorption from aqueur Phase by activited Corbon: a simplified application of the Salonphobia Newy" X, Xirmalakhorley & R. C. Speece

See 9-75. *

Polorographic analyses of advolving power for He & Blue and H30 t of HCl
"decolocation of HB d.n.g analos mm ms of e alspu P-1 t-il royac ionol.
288
Henry 28: 405 Berlawan 5 51347 Elian H PS 47 Seager CD 46 Known and 18 50 The A MRP 47
Tylu A ARP 47
See Eyster 1943 JCCP
Exted by Boding St '50 phanol 42: 78:45-40
K Henry 13 13 July 1 19
Melman # Browlef PS Sevay MG Y6 Adv 95 JB
(sundan M 51

Gosman BA Chimie & beduttie Spec No. 199-203 (1933 pm) Study of theadrochent power of extinated charcoal ky the polarographic method.

rites Eyster 8/98

HENRY MD—THE EFFECT OF PENICILLIN ON METHYLENE BLUE ADSORPTION ONTO ACTIVATED CHARCOAL J GEN PHYSL 28(5):415-419

Institute for Scientific Information®

CITATION DATABASES













Cited References--Full Record

Article 1 of 5





RELATED RECORDS

ates tyslu

Studies on the adsorption of sulfapyridine at the solution-alumina interface Bajpai AK, Rajpoot M, Mishra DD JOURNAL OF COLLOID AND INTERFACE SCIENCE

187: (1) 96-104 MAR 1 1997

Document type: Article Language: English Cited References: 31 Times Cited: 1

Abstract:

The adsorption of Sulfapyridine (SP) onto the alumina surface has been carried out at room temperature to study the adsorption behavior of SP and its mode of interaction with the surface of alumina particles. It is found that various factors such as concentration of SP solution, required time for adsorption equilibrium, pH of the adsorbate solution, temperature of adsorption medium, presence of ions like PO43-, SO42-, and Cl- affect quantitatively the adsorption of SP. Various adsorption and kinetic parameters such as the adsorption isotherm, adsorption coefficient, rate constants for adsorption and desorption and surface coverage have also been evaluated. The present study is significant because knowledge of the exact interaction between the SP molecules and the alumina surface and the proper choice of experimental condition would be helpful in carrying out an optimum separation of sulfa drug compounds chromatographically. (C) 1997 Academic Press.

Author Keywords:

sulfapyridine, adsorption, kinetics

KeyWords Plus:

KINETICS, SURFACES

Addresses:

Bajpai AK, GOVT AUTOMOMOUS SCI COLL, DEPT CHEM, BOSE MEM RES LAB, JABALPUR 482001, MADHYA PRADESH, INDIA.

RANI DURGAVATI UNIV, DEPT POSTGRAD STUDIES & RES CHEM, JABALPUR 482001, MADHYA PRADESH, INDIA.

Publisher:

ACADEMIC PRESS INC JNL-COMP SUBSCRIPTIONS, SAN DIEGO

IDS Number:

WN142

ISSN:

0021-9797